

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

MHC II (HLA-DRA)(19-26.1), CF568 conjugate, 0.1mg/mL, Clone: [19-26.1], Mouse, Monoclonal BOT-BNC681082-100

| | |
|----------------------------|--|
| Article Name | MHC II (HLA-DRA)(19-26.1), CF568 conjugate, 0.1mg/mL, Clone: [19-26.1], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC681082-100 |
| Supplier Catalog Number | BNC681082-100 |
| Alternative Catalog Number | BOT-BNC681082-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | FC, IHC |
| Species Reactivity | Human |
| Immunogen | Raji cells |
| Conjugation | CF568 |
| Product Description | This MAbs reacts with the HLA-DR antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36 kDa alpha (heavy) chain and a 28 kDa beta (light) chain... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [19-26.1] |

| | |
|-------------------|---|
| Molecular Weight | 36 kDa (? chain) and 27 kDa (? chain) |
| UniProt | P01903 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |