

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Ep-CAM / CD326 (Epithelial Marker) (EGP40/1555R), CF568 conjugate, 0.1mg/mL, Clone: [EGP40/1555R], Rabbit, Monoclonal BOT-BNC681555-500**

Article Name	Ep-CAM / CD326 (Epithelial Marker) (EGP40/1555R), CF568 conjugate, 0.1mg/mL, Clone: [EGP40/1555R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNC681555-500
Supplier Catalog Number	BNC681555-500
Alternative Catalog Number	BOT-BNC681555-500-500UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	FC, IHC
Species Reactivity	Human
Immunogen	A synthetic peptide (around aa 20-60) from the N-terminus of human TACSTD1 protein (exact sequence is proprietary)
Conjugation	CF568
Product Description	This antibody recognizes a 40-43 kDa transmembrane epithelial glycoprotein, identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). Ep-CAM is expressed on baso-lateral cell surface in most simple epithelia ...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[EGP40/1555R]

Molecular Weight	40-43 kDa
UniProt	<a href="#">P16422</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user