

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

FOXA1 / HNF3A (FOXA1/2230R), CF568 conjugate, 0.1mg/mL, Clone: [FOXA1/2230R], Rabbit, Monoclonal BOT-BNC682230-500

| | |
|----------------------------|---|
| Article Name | FOXA1 / HNF3A (FOXA1/2230R), CF568 conjugate, 0.1mg/mL, Clone: [FOXA1/2230R], Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNC682230-500 |
| Supplier Catalog Number | BNC682230-500 |
| Alternative Catalog Number | BOT-BNC682230-500-500UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | FC, IF, IHC, WB |
| Species Reactivity | Human, Rat |
| Immunogen | Recombinant human FOXA1 protein fragment (around aa372-472) (exact sequence is proprietary) |
| Conjugation | CF568 |
| Product Description | The transcription factor Forkhead-box A1 (FOXA1), also known as hepatocyte nuclear factor 3-alpha, is a member of the FOX class of transcription factors. HNF-1 (alpha and beta), HNF-3 (alpha, beta and gamma), HNF-4 (alpha and gamma), and HNF-6 compos... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [FOXA1/2230R] |

| | |
|-------------------|--|
| Molecular Weight | 49 kDa |
| UniProt | P55317 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |