

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

HCG-beta(HCGb/54), CF488A conjugate, 0.1mg/mL, Clone: [HCGb/54], Mouse, Monoclonal BOT-BNC880054-500

| | |
|----------------------------|---|
| Article Name | HCG-beta(HCGb/54), CF488A conjugate, 0.1mg/mL, Clone: [HCGb/54], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC880054-500 |
| Supplier Catalog Number | BNC880054-500 |
| Alternative Catalog Number | BOT-BNC880054-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant hCG beta protein |
| Conjugation | CF488A |
| Product Description | This MAb reacts with a protein of 22 kDa, identified as beta sub-unit of HCG. It does not cross react with the alpha sub-unit. HCG is a glycoprotein, which is secreted in large quantities by normal trophoblasts. It is present only in trace amounts in... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [HCGb/54] |

| | |
|-------------------|---|
| Molecular Weight | 22 kDa |
| UniProt | P01233 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |