

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Granulocyte Marker(BM-1), CF488A conjugate, 0.1mg/mL, IgG1, Clone: [BM-1], Mouse, Monoclonal BOT-BNC880061-500

| | |
|----------------------------|--|
| Article Name | Granulocyte Marker(BM-1), CF488A conjugate, 0.1mg/mL, IgG1, Clone: [BM-1], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNC880061-500 |
| Supplier Catalog Number | BNC880061-500 |
| Alternative Catalog Number | BOT-BNC880061-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IF, IHC, RIA, WB |
| Species Reactivity | Human |
| Immunogen | Human peripheral blood mononuclear cells |
| Conjugation | CF488A |
| Product Description | Recognizes 183 kDa protein with DNA-binding characteristics, which is identified as a myeloid specific antigen. It reacts with myeloid precursor cells and granulocytes in bone marrow. Its antigen appears to be restricted to M2 and M3 acute myelogenous... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [BM-1] |

| | |
|-------------------|--|
| Molecular Weight | 183 kDa |
| Isotype | IgG1 |
| UniProt | Not Known |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL No special pretreatment is required for staining of formalin/paraffin tissues Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |