

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **PCNA(PC10), CF594 conjugate, 0.1mg/mL, Clone: [PC10], Mouse, Monoclonal BOT-BNC940467-500**

Article Name	PCNA(PC10), CF594 conjugate, 0.1mg/mL, Clone: [PC10], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNC940467-500
Supplier Catalog Number	BNC940467-500
Alternative Catalog Number	BOT-BNC940467-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	FC, IHC
Species Reactivity	Drosophila, Gallus, Human, Monkey, Mouse, Porcine, Rat, Yeast, Zebrafish
Immunogen	Rat PCNA/Protein A fusion protein
Conjugation	CF594
Product Description	Recognizes a non-histone protein of 36 kDa, which is identified as proliferating cell nuclear antigen (PCNA). It is also known as cyclin or polymerase delta auxiliary protein. Elevated expression of PCNA/cyclin has been shown in the nucleus during la...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[PC10]

Molecular Weight	36 kDa
UniProt	<a href="#">P12004</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user