

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Mucin 1 / EMA / Episialin / CD227(GP1.4), Biotin conjugate, 0.1mg/mL, Clone: [GP1.4], Mouse, Monoclonal BOT-BNCB0159-500**

Article Name	Mucin 1 / EMA / Episialin / CD227(GP1.4), Biotin conjugate, 0.1mg/mL, Clone: [GP1.4], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNCB0159-500
Supplier Catalog Number	BNCB0159-500
Alternative Catalog Number	BOT-BNCB0159-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC, WB
Species Reactivity	Human
Immunogen	Human milk fat globule membranes
Conjugation	Biotin
Product Description	In Western blotting, this antibody recognizes proteins in MW range of 265-400 kDa, identified as different glycoforms of EMA. This MAb reacts with the DTRP epitope in the tandem repeats. The alpha subunit has cell adhesive properties. It can act both ...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[GP1.4]

Molecular Weight	265-400 kDa
UniProt	<a href="#">P15941</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.1-0.2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user