

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

p40(Rabbit PAb), Biotin conjugate, 0.1mg/mL, Clone: [PAb], Monoclonal BOT-BNCB0375-500

| | |
|----------------------------|---|
| Article Name | p40(Rabbit PAb), Biotin conjugate, 0.1mg/mL, Clone: [PAb], Monoclonal |
| Biozol Catalog Number | BOT-BNCB0375-500 |
| Supplier Catalog Number | BNCB0375-500 |
| Alternative Catalog Number | BOT-BNCB0375-500-500UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Bovine, Human, Mouse, Rat |
| Immunogen | A synthetic peptide (ENNAQTQFSEPQY) corresponding to aa5-17 of human p40 |
| Conjugation | Biotin |
| Product Description | p40 (p63 delta) is a marker determined to be highly specific for squamous basal cells by immunohistochemistry (IHC). The current more routinely recommended marker, p63, appears to have less specificity compared to p40, especially on squamous cell tum... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [PAb] |

| | |
|-------------------|--|
| Molecular Weight | 40 kDa |
| UniProt | Q9H3D4 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |