

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

BCL-10(BL10/411), Biotin conjugate, 0.1mg/mL, Clone: [BL10/411], Mouse, Monoclonal BOT-BNCB0411-500

Article Name	BCL-10(BL10/411), Biotin conjugate, 0.1mg/mL, Clone: [BL10/411], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNCB0411-500
Supplier Catalog Number	BNCB0411-500
Alternative Catalog Number	BOT-BNCB0411-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	FC, IHC, WB
Species Reactivity	Human
Immunogen	Human BCL10 recombinant protein (epitope aa122-168)
Conjugation	Biotin
Product Description	B-cell lymphoma/leukemia 10 (BCL10) is a signaling protein that promotes apoptosis, pro-caspase-9 maturation, and activation of NF-kappa-B via NIK and IKK. BCL10 contains an N-terminal caspase recruitment domain (CARD), which is found in a number of ...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[BL10/411]

Molecular Weight	33 kDa
UniProt	O95999
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user