

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Mucin 1 / EMA / Episialin / CD227(E29), Biotin conjugate, 0.1mg/mL, Clone: [E29], Mouse, Monoclonal BOT-BNCB0478-100**

Article Name	Mucin 1 / EMA / Episialin / CD227(E29), Biotin conjugate, 0.1mg/mL, Clone: [E29], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNCB0478-100
Supplier Catalog Number	BNCB0478-100
Alternative Catalog Number	BOT-BNCB0478-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Delipidated extract of human milk fat globule membranes
Conjugation	Biotin
Product Description	In Western blotting, this antibody recognizes proteins in MW range of 265-400 kDa, identified as different glycoforms of EMA. EMA may provide a protective layer on epithelial cells against bacterial and enzyme attack. In immunohistochemical assays, i...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[E29]

Molecular Weight	265-400 kDa
UniProt	<a href="#">P15941</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.1-0.2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user