

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Moesin(MSN/492), Biotin conjugate, 0.1mg/mL, Clone: [MSN/492], Mouse, Monoclonal BOT-BNCB0492-500**

Article Name	Moesin(MSN/492), Biotin conjugate, 0.1mg/mL, Clone: [MSN/492], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNCB0492-500
Supplier Catalog Number	BNCB0492-500
Alternative Catalog Number	BOT-BNCB0492-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	FC, IF, IHC, WB
Species Reactivity	Human
Immunogen	Recombinant full-length human Moesin protein
Conjugation	Biotin
Product Description	Recognizes 78 kDa moesin protein. Moesin, a member of the talin-4.1 superfamily, is a linking protein of the sub-membranous actin cytoskeleton. It is expressed in variable amounts in cells of different phenotypes such as macrophages, lymphocytes, fib...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[MSN/492]

Molecular Weight	78 kDa
UniProt	<a href="#">P26038</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Does not react with rat, others not known Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user