

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Alkaline Phosphatase (Placental) / PLAP (Germ Cell Tumor Marker)(ALPP/516), Biotin conjugate, 0.1mg/mL, Clone: [ALPP/516], Mouse, Monoclonal BOT-BNCB0516-500

| | |
|----------------------------|---|
| Article Name | Alkaline Phosphatase (Placental) / PLAP (Germ Cell Tumor Marker)(ALPP/516), Biotin conjugate, 0.1mg/mL, Clone: [ALPP/516], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNCB0516-500 |
| Supplier Catalog Number | BNCB0516-500 |
| Alternative Catalog Number | BOT-BNCB0516-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human ALPP protein |
| Conjugation | Biotin |
| Product Description | This antibody reacts with placental alkaline phosphatase. There are at least four distinct but related alkaline phosphatases: intestinal, placental (PLAP), placental-like, and liver/bone/kidney (tissue non-specific). The first three are located toget... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [ALPP/516] |

| | |
|-------------------|--|
| Molecular Weight | 70 kDa |
| UniProt | P05187 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |