

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Mucin 1 / EMA / Episialin / CD227(MUC1/520), Biotin conjugate, 0.1mg/mL, Clone: [MUC1/520], Mouse, Monoclonal BOT-BNCB0520-100

Article Name	Mucin 1 / EMA / Episialin / CD227(MUC1/520), Biotin conjugate, 0.1mg/mL, Clone: [MUC1/520], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNCB0520-100
Supplier Catalog Number	BNCB0520-100
Alternative Catalog Number	BOT-BNCB0520-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC, WB
Species Reactivity	Human
Immunogen	Human milk-fat globule membranes (HMFGM)
Conjugation	Biotin
Product Description	In Western blotting, it recognizes proteins in MW range of 265-400 kDa, identified as different glycoforms of EMA. EMA may provide a protective layer on epithelial cells against bacterial and enzyme attack. In immunohistochemical assays, it superbly ...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[MUC1/520]

Molecular Weight	265-400 kDa
UniProt	P15941
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.1-0.2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user