

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker)(Ber-H2), Biotin conjugate, 0.1mg/mL, Clone: [Ber-H2], Mouse, Monoclonal BOT-BNCB0772-500

Article Name	CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker)(Ber-H2), Biotin conjugate, 0.1mg/mL, Clone: [Ber-H2], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNCB0772-500
Supplier Catalog Number	BNCB0772-500
Alternative Catalog Number	BOT-BNCB0772-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Species Reactivity	Human
Immunogen	Co cell line established from a patient with Hodgkin's disease of T-cell lineage
Conjugation	Biotin
Product Description	This antibody recognizes a single chain glycoprotein of 105/120 kDa, identified as CD30/Ki-1. Its epitope is located between aa112-412. CD30 is synthesized as a 90 kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphory...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[Ber-H2]
Molecular Weight	105-120 kDa

UniProt	P28908
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry