

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Golgi Complex(AE-6), Biotin conjugate, 0.1mg/mL, Clone: [AE-6], Mouse, Monoclonal BOT-BNCB0868-100

| | |
|----------------------------|---|
| Article Name | Golgi Complex(AE-6), Biotin conjugate, 0.1mg/mL, Clone: [AE-6], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNCB0868-100 |
| Supplier Catalog Number | BNCB0868-100 |
| Alternative Catalog Number | BOT-BNCB0868-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | SU-DHL-1 large cell lymphoma cells. |
| Conjugation | Biotin |
| Product Description | This MAb recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining ... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [AE-6] |

| | |
|-------------------|---|
| Molecular Weight | Not Known |
| UniProt | Not Known |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Does not react with mouse or rat, others not known Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Immunocytochemistry Acetone or paraformaldehyde fixed 0.5-1 ug/mL Western blotting 0.5-1.0 ug/mL Flow Cytometry 0.5-1.0 ug/million cells in 0.1 mL Optimal dilution for a specific application should be determined by user |