

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD54(W-CAM-1, same as Wehi-CAM-1 or 1H4), Biotin conjugate, 0.1mg/mL, Clone: [W-CAM-1, same as Wehi-CAM-1 or 1H4], Mouse, Monoclonal BOT-BNCB0988-100

| | |
|----------------------------|---|
| Article Name | CD54(W-CAM-1, same as Wehi-CAM-1 or 1H4), Biotin conjugate, 0.1mg/mL, Clone: [W-CAM-1, same as Wehi-CAM-1 or 1H4], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNCB0988-100 |
| Supplier Catalog Number | BNCB0988-100 |
| Alternative Catalog Number | BOT-BNCB0988-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Raji Burkitt lymphoma cells |
| Conjugation | Biotin |
| Product Description | Recognizes an 85-115 kDa protein (variation with cell type), identified as intercellular adhesion molecule (ICAM-1) (Workshop IV). It has 7 potential N-linked glycosylation sites. ICAM-1 is a single chain glycoprotein of Ig supergene family, present ... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [W-CAM-1, same as Wehi-CAM-1 or 1H4] |

| | |
|-------------------|---|
| Molecular Weight | 85-115 kDa |
| UniProt | P05362 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Immunohistology formalin-fixed 2-4 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Immunofluorescence 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Functional Studies order Ab without BSA & azide) Optimal dilution for a specific application should be determined by user |