

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin, pan (Cocktail PAN-CK), Biotin conjugate, 0.1mg/mL, Clone: [Cocktail PAN-CK], Mouse, Monoclonal BOT-BNCB0999-100

Article Name	Cytokeratin, pan (Cocktail PAN-CK), Biotin conjugate, 0.1mg/mL, Clone: [Cocktail PAN-CK], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNCB0999-100
Supplier Catalog Number	BNCB0999-100
Alternative Catalog Number	BOT-BNCB0999-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	FC, IF, IHC
Species Reactivity	Bovine, Canine, Gallus, Human, Monkey, Mouse, Rabbit, Rat
Immunogen	Human epidermal keratin
Conjugation	Biotin
Product Description	Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subfamilies. This antibody cocktail recognizes acidic (Type I or LMW) and basic (Type II or HMW) cytokeratins, with 67 kDa (CK1), 64 kDa (CK3), 59 kDa (C...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[Cocktail PAN-CK]

Molecular Weight	40-67 kDa (Multiple)
UniProt	Q7Z794
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user