

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Complement C4d(TDM16), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [TDM16], Mouse, Monoclonal BOT-BNCB1232-500

| | |
|----------------------------|---|
| Article Name | Complement C4d(TDM16), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [TDM16], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNCB1232-500 |
| Supplier Catalog Number | BNCB1232-500 |
| Alternative Catalog Number | BOT-BNCB1232-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Species Reactivity | Human |
| Immunogen | Human Complement 4d protein |
| Conjugation | Biotin |
| Product Description | This MAb is specific to Complement 4d (C4d) and it reacts with the secreted as well as cell-bound C4d.C4d is a degradation product of the activated complement factor C4b. Complement 4b is typically activated by binding of Abs to specific target molec... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [TDM16] |
| Molecular Weight | 192 kDa (predicted) |

| | |
|-------------------|--|
| Isotype | IgG1 |
| UniProt | P0COL4 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |