

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

TDP2 / EAPII (TDP2/1258), Biotin conjugate, 0.1mg/mL, Clone: [TDP2/1258], Mouse, Monoclonal BOT-BNCB1258-500

| | |
|----------------------------|---|
| Article Name | TDP2 / EAPII (TDP2/1258), Biotin conjugate, 0.1mg/mL, Clone: [TDP2/1258], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNCB1258-500 |
| Supplier Catalog Number | BNCB1258-500 |
| Alternative Catalog Number | BOT-BNCB1258-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant human full-length TDP2 protein |
| Conjugation | Biotin |
| Product Description | This MAb recognizes a protein of 41 kDa, which is identified as TDP2. It is a member of a superfamily of divalent cation-dependent phosphodiesterases. The encoded protein associates with CD40, tumor necrosis factor (TNF) receptor-75 and TNF receptor ... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [TDP2/1258] |

| | |
|-------------------|--|
| Molecular Weight | ~41 kDa |
| UniProt | O95551 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |