

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Caldesmon, HMW (h-Caldesmon) (Smooth Muscle Marker) (CALD1/1424R), Biotin conjugate, 0.1mg/mL, Clone: [CALD1/1424R], Rabbit, Monoclonal BOT-BNCB1424-500**

|                            |   |
|----------------------------|---|
| Article Name               | Caldesmon, HMW (h-Caldesmon) (Smooth Muscle Marker) (CALD1/1424R), Biotin conjugate, 0.1mg/mL, Clone: [CALD1/1424R], Rabbit, Monoclonal   |
| Biozol Catalog Number      | BOT-BNCB1424-500  |
| Supplier Catalog Number    | BNCB1424-500  |
| Alternative Catalog Number | BOT-BNCB1424-500-500UL  |
| Manufacturer               | Biotium   |
| Host                       | Rabbit  |
| Category                   | Antikörper  |
| Application                | IHC, WB   |
| Species Reactivity         | Human, Rat  |
| Immunogen                  | Recombinant human full-length CALD1 protein   |
| Conjugation                | Biotin  |
| Product Description        | Caldesmon HMW is the high molecular weight variant of Caldesmon. Two closely related variants of human caldesmon have been identified which are different in their electrophoretic mobility and cellular distribution. The h-caldesmon variant (120-150 kD... |
| Clonality                  | Monoclonal  |
| Concentration              | 0.1 mg/mL   |
| Clone Designation          | [CALD1/1424R]   |

|                   |  |
|-------------------|--|
| Molecular Weight  | 150 kDa  |
| UniProt           | <a href="#">Q05682</a>   |
| Buffer            | PBS, 0.1% BSA, 0.05% azide   |
| Source            | Animal   |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 1 mM EDTA pH 7.5-8.5 Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |