

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD79a (IGA/1688R), Biotin conjugate, 0.1mg/mL, Clone: [IGA/1688R], Rabbit, Monoclonal BOT-BNCB1688-100

| | |
|----------------------------|---|
| Article Name | CD79a (IGA/1688R), Biotin conjugate, 0.1mg/mL, Clone: [IGA/1688R], Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNCB1688-100 |
| Supplier Catalog Number | BNCB1688-100 |
| Alternative Catalog Number | BOT-BNCB1688-100-100UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | FC, IF, IHC, WB |
| Species Reactivity | Human, Mouse, Rat |
| Immunogen | A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79a protein. |
| Conjugation | Biotin |
| Product Description | A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [IGA/1688R] |

| | |
|-------------------|--|
| Molecular Weight | 44 kDa |
| UniProt | P11912 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |