

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Bcl-X (Apoptosis Marker)(BCL2L1/1762), Biotin conjugate, 0.1mg/mL, Clone: [BCL2L1/1762], Mouse, Monoclonal BOT-BNCB1762-500

Article Name	Bcl-X (Apoptosis Marker)(BCL2L1/1762), Biotin conjugate, 0.1mg/mL, Clone: [BCL2L1/1762], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNCB1762-500
Supplier Catalog Number	BNCB1762-500
Alternative Catalog Number	BOT-BNCB1762-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Species Reactivity	Human
Immunogen	Recombinant human BCL2L1 protein fragment (around aa 1-135) (exact sequence is proprietary)
Conjugation	Biotin
Product Description	Bcl-x belongs to the Bcl-2 family of proteins and plays a crucial role in apoptosis. Bcl-X has two isoforms: Bcl-XL (long), a 26 kDa protein which suppresses cell death, and Bcl-XS (short), a 178 amino acid protein lacking a 63 amino acid domain whic...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[BCL2L1/1762]
Molecular Weight	27 kDa

UniProt	Q07817
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry