

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) (VWF/1767), Biotin conjugate, 0.1mg/mL, Clone: [VWF/1767], Mouse, Monoclonal BOT-BNCB1767-500

| | |
|----------------------------|---|
| Article Name | von Willebrand Factor / Factor VIII Related-Ag (Endothelial Marker) (VWF/1767), Biotin conjugate, 0.1mg/mL, Clone: [VWF/1767], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNCB1767-500 |
| Supplier Catalog Number | BNCB1767-500 |
| Alternative Catalog Number | BOT-BNCB1767-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment of human vWF protein (aa1815-1939) (exact sequence is proprietary) |
| Conjugation | Biotin |
| Product Description | von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslation... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |

| | |
|-------------------|--|
| Clone Designation | [VWF/1767] |
| Molecular Weight | 250 kDa |
| UniProt | P04275 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/mL for coating order Ab without BSA Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |