

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Carcinoembryonic Antigen (CEA) / CD66 (C66/2055R), Biotin conjugate, 0.1mg/mL, Clone: [C66/2055R], Rabbit, Monoclonal BOT-BNCB2055-100

| | |
|----------------------------|---|
| Article Name | Carcinoembryonic Antigen (CEA) / CD66 (C66/2055R), Biotin conjugate, 0.1mg/mL, Clone: [C66/2055R], Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNCB2055-100 |
| Supplier Catalog Number | BNCB2055-100 |
| Alternative Catalog Number | BOT-BNCB2055-100-100UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human CEA protein |
| Conjugation | Biotin |
| Product Description | This antibody recognizes proteins of 80-200 kDa, identified as different members of CEA family. CEA is synthesized during development in the fetal gut and is re-expressed in increased amounts in intestinal carcinomas and several other tumors. This MA... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [C66/2055R] |

| | |
|-------------------|---|
| Molecular Weight | 80-200 kDa |
| UniProt | P06731 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |