

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **TCL1 (T-Cell Marker) (TCL1/2078), Biotin conjugate, 0.1mg/mL, Clone: [TCL1/2078], Mouse, Monoclonal BOT-BNCB2078-100**

Article Name	TCL1 (T-Cell Marker) (TCL1/2078), Biotin conjugate, 0.1mg/mL, Clone: [TCL1/2078], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNCB2078-100
Supplier Catalog Number	BNCB2078-100
Alternative Catalog Number	BOT-BNCB2078-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant human TCL1 protein fragment (around aa 2-109) (exact sequence is proprietary)
Conjugation	Biotin
Product Description	T-cell leukemia/Lymphoma Protein 1A (TCL-1A), also known as p14TCL1 is a product of the TCL1 gene that is involved in T-cell prolymphocytic leukemia (TPLL). T-PLL is a rare form of mature T-cell leukemia, which is consistently associated with chromos...
Clonality	Monoclonal
Concentration	0.1 mg/mL
Clone Designation	[TCL1/2078]

Molecular Weight	14 kDa
UniProt	<a href="#">P56279</a>
Buffer	PBS, 0.1% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user