

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD52 (Epididymis-Specific Protein 5) (CD52/2276R), Biotin conjugate, 0.1mg/mL, Clone: [CD52/2276R], Rabbit, Monoclonal BOT-BNCB2276-500

| | |
|----------------------------|--|
| Article Name | CD52 (Epididymis-Specific Protein 5) (CD52/2276R), Biotin conjugate, 0.1mg/mL, Clone: [CD52/2276R], Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNCB2276-500 |
| Supplier Catalog Number | BNCB2276-500 |
| Alternative Catalog Number | BOT-BNCB2276-500-500UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human, Monkey |
| Immunogen | Recombinant full-length human CD52 protein |
| Conjugation | Biotin |
| Product Description | Human CD52 is expressed at high density by lymphocytes, monocytes, eosinophils, thymocytes and macrophages. It is expressed by most lymphoid derived malignancies, although expression on myeloma cells is variable. Primary antibodies are available purif... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [CD52/2276R] |

| | |
|-------------------|---|
| Molecular Weight | 6 kDa |
| UniProt | P31358 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |