

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD34 (Hematopoietic Stem Cell & Endothelial Marker) (HPCA1/2598R), Biotin conjugate, 0.1mg/mL, Clone: [HPCA1/2598R], Rabbit, Monoclonal BOT-BNCB2598-100

| | |
|----------------------------|---|
| Article Name | CD34 (Hematopoietic Stem Cell & Endothelial Marker) (HPCA1/2598R), Biotin conjugate, 0.1mg/mL, Clone: [HPCA1/2598R], Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNCB2598-100 |
| Supplier Catalog Number | BNCB2598-100 |
| Alternative Catalog Number | BOT-BNCB2598-100-100UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | FC, IHC |
| Species Reactivity | Human, Rat |
| Immunogen | Recombinant full-length human HPCA1 protein |
| Conjugation | Biotin |
| Product Description | This antibody recognizes a transmembrane, heavily glycosylated protein of 90-120 kDa, which is identified as CD34. Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as li... |
| Clonality | Monoclonal |
| Concentration | 0.1 mg/mL |
| Clone Designation | [HPCA1/2598R] |

| | |
|-------------------|--|
| Molecular Weight | 90-110 kDa |
| UniProt | P28906 |
| Buffer | PBS, 0.1% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |