

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **HCG-intact(HCGab/52), 0.2mg/mL, Clone: [HCGab/52], Mouse, Monoclonal BOT-BNUB0052-500**

Article Name	HCG-intact(HCGab/52), 0.2mg/mL, Clone: [HCGab/52], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB0052-500
Supplier Catalog Number	BNUB0052-500
Alternative Catalog Number	BOT-BNUB0052-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Purified hCG protein
Product Description	This MAb is a very special because it reacts ONLY with the intact-HCG and not with either free alpha- or free beta-chain of HCG. HCG is a glycoprotein and is composed of two non-identical, non-covalently linked polypeptide chains designated as the al...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[HCGab/52]
Molecular Weight	~35 kDa (intact hCG)
UniProt	<a href="#">P01215</a>

Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user