

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

c-Myc(MYC275), 0.2mg/mL, Clone: [MYC275], Mouse, Monoclonal BOT-BNUB0275-500

| | |
|----------------------------|---|
| Article Name | c-Myc(MYC275), 0.2mg/mL, Clone: [MYC275], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNUB0275-500 |
| Supplier Catalog Number | BNUB0275-500 |
| Alternative Catalog Number | BOT-BNUB0275-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant human c-myc protein |
| Product Description | The c-Myc protein is a transcription factor, which is encoded by the c-Myc gene on human chromosome 8q24. c-Myc is commonly activated in a variety of tumor cells and plays an important role in cellular proliferation, differentiation, apoptosis and ce... |
| Clonality | Monoclonal |
| Concentration | 0.2 mg/mL |
| Clone Designation | [MYC275] |
| Molecular Weight | 62-64 kDa |
| UniProt | P01106 |

| | |
|-------------------|--|
| Buffer | PBS, 0.05% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user |