

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

NKX2.2(NX2/294), 0.2mg/mL, Clone: [NX2/294], Mouse, Monoclonal BOT-BNUB0294-500

| | |
|----------------------------|---|
| Article Name | NKX2.2(NX2/294), 0.2mg/mL, Clone: [NX2/294], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNUB0294-500 |
| Supplier Catalog Number | BNUB0294-500 |
| Alternative Catalog Number | BOT-BNUB0294-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Gallus, Human, Mouse, Rat |
| Immunogen | Human full-length recombinant NKX2.2 protein |
| Product Description | Expression of NKX2.2 has been found in neuroendocrine tumors of the gut, making it a potential marker for the study of gastrointestinal neuroendocrine tumors. More recently, NKX2.2 protein was identified as a target of EWS-FLI-1, the fusion protein s... |
| Clonality | Monoclonal |
| Concentration | 0.2 mg/mL |
| Clone Designation | [NX2/294] |
| Molecular Weight | 40-50 kDa |
| UniProt | O95096 |

| | |
|-------------------|--|
| Buffer | PBS, 0.05% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |