

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

TNF alpha(TNF706 + P/T2), 0.2mg/mL, Clone: [TNF706], Mouse, Monoclonal BOT-BNUB0707-100

| | |
|----------------------------|---|
| Article Name | TNF alpha(TNF706 + P/T2), 0.2mg/mL, Clone: [TNF706], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNUB0707-100 |
| Supplier Catalog Number | BNUB0707-100 |
| Alternative Catalog Number | BOT-BNUB0707-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IF, IHC |
| Species Reactivity | Canine, Feline, Human, Mouse, Rabbit, Rat, Zebrafish |
| Immunogen | Recombinant N-terminal fragment of human TNF-? |
| Product Description | TNF alpha is a protein secreted by lipopolysaccharide-stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF alpha exists as a multimer of two, three, or five non-covalently linked units, but shows a single 17 kD... |
| Clonality | Monoclonal |
| Concentration | 0.2 mg/mL |
| Clone Designation | [TNF706] |
| Molecular Weight | 17 kDa |
| UniProt | P01375 |

| | |
|-------------------|--|
| Buffer | PBS, 0.05% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Immunohistology formalin-paraffin 2-4 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris buffer with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user |