

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cytokeratin 17(KRT17/778), 0.2mg/mL, Clone: [KRT17/778], Mouse, Monoclonal BOT-BNUB0778-100

Article Name	Cytokeratin 17(KRT17/778), 0.2mg/mL, Clone: [KRT17/778], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB0778-100
Supplier Catalog Number	BNUB0778-100
Alternative Catalog Number	BOT-BNUB0778-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Bovine, Goat, Human, Porcine, Rat
Immunogen	Recombinant full-length human KRT17 protein
Product Description	Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excellent tool to distinguish myoepithelial cells from luminal epithelium of various glands such as ...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[KRT17/778]
Molecular Weight	46 kDa

UniProt	Q04695
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user