

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

TGFalpha(TG86), 0.2mg/mL, Clone: [TG86], Mouse, Monoclonal BOT-BNUB0786-500

| | |
|----------------------------|---|
| Article Name | TGFalpha(TG86), 0.2mg/mL, Clone: [TG86], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNUB0786-500 |
| Supplier Catalog Number | BNUB0786-500 |
| Alternative Catalog Number | BOT-BNUB0786-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human, Rabbit, Zebrafish |
| Immunogen | A 10-amino acid synthetic peptide (aa34-43, PPVAAAVVSH) from human TGF α . |
| Product Description | This antibody reacts with the TGF alpha and shows no cross-reaction with EGF and the neuropeptide synenkephalin. The staining with this MAb is completely blocked by the peptide used for raising this antibody. TGF-beta (aa50) is a growth factor with 3... |
| Clonality | Monoclonal |
| Concentration | 0.2 mg/mL |
| Clone Designation | [TG86] |
| Molecular Weight | 6 kDa |
| UniProt | P01135 |

| | |
|-------------------|--|
| Buffer | PBS, 0.05% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Immunohistology formalin-paraffin 2-4 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user |