

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Ep-CAM / CD326(EGP40/837), 0.2mg/mL, Clone: [EGP40/837], Mouse, Monoclonal BOT-BNUB0837-100

Article Name	Ep-CAM / CD326(EGP40/837), 0.2mg/mL, Clone: [EGP40/837], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB0837-100
Supplier Catalog Number	BNUB0837-100
Alternative Catalog Number	BOT-BNUB0837-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	FC, IF, IHC, WB
Species Reactivity	Human
Immunogen	Recombinant human TACSTD1 protein
Product Description	EGP40 is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majo...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[EGP40/837]
Molecular Weight	40-43 kDa

UniProt	P16422
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not react with mouse or rat, others not known Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user