

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker)(Ber-H2 + CD30/412), 0.2mg/mL, Clone: [Ber-H2 CD30/412], Mouse, Monoclonal BOT-BNUB0849-100

|                            |   |
|----------------------------|---|
| Article Name               | CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker)(Ber-H2 + CD30/412), 0.2mg/mL, Clone: [Ber-H2 CD30/412], Mouse, Monoclonal   |
| Biozol Catalog Number      | BOT-BNUB0849-100  |
| Supplier Catalog Number    | BNUB0849-100  |
| Alternative Catalog Number | BOT-BNUB0849-100-100UL  |
| Manufacturer               | Biotium   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Species Reactivity         | Human   |
| Immunogen                  | Co cell line established from a patient with Hodgkin's disease of T-cell lineage (Ber-H2), human CD30 recombinant protein (CD30/412)  |
| Product Description        | This antibody recognizes a single chain glycoprotein of 105/120 kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90 kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120 kDa glycoprotein. In ... |
| Clonality                  | Monoclonal  |
| Concentration              | 0.2 mg/mL   |
| Clone Designation          | [Ber-H2 CD30/412]   |
| Molecular Weight           | 105-120 kDa   |

|                   |   |
|-------------------|---|
| UniProt           | P28908  |
| Buffer            | PBS, 0.05% BSA, 0.05% azide   |
| Source            | Animal  |
| Application Notes | <p>For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry</p> |