

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Golgi Complex(AE-6), 0.2mg/mL, Clone: [AE-6], Mouse, Monoclonal BOT-BNUB0868-500**

Article Name	Golgi Complex(AE-6), 0.2mg/mL, Clone: [AE-6], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB0868-500
Supplier Catalog Number	BNUB0868-500
Alternative Catalog Number	BOT-BNUB0868-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	SU-DHL-1 large cell lymphoma cells.
Product Description	This MAb recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining ...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[AE-6]
Molecular Weight	Not Known
UniProt	Not Known

Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Does not react with mouse or rat, others not known Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Immunocytochemistry Acetone or paraformaldehyde fixed 0.5-1 ug/mL Western blotting 0.5-1.0 ug/mL Flow Cytometry 0.5-1.0 ug/million cells in 0.1 mL Optimal dilution for a specific application should be determined by user</p>