

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **CD99 / MIC2 (Ewings Sarcoma Marker)(12E7), 0.2mg/mL, Clone: [1.20E 08], Mouse, Monoclonal BOT-BNUB0876-100**

Article Name	CD99 / MIC2 (Ewings Sarcoma Marker)(12E7), 0.2mg/mL, Clone: [1.20E 08], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB0876-100
Supplier Catalog Number	BNUB0876-100
Alternative Catalog Number	BOT-BNUB0876-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Species Reactivity	Human
Immunogen	Human acute lymphocytic leukemia T-cells
Product Description	This antibody recognizes a sialoglycoprotein of 27-32 kDa, identified as CD99, or MIC2 gene product, or E2 antigen. MIC2 gene is located in the pseudo-autosomal region of the human X and Y chromosome. MIC2 gene encodes two distinct proteins, which ar...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[1.20E 08]
Molecular Weight	27-32 kDa
UniProt	<a href="#">P14209</a>

Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry