

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Thomsen-Friedenreich Antigen / CD176 (Pan Carcinoma Marker)(A78-G/A7), 0.2mg/mL, Clone: [A78-G/A7], Mouse, Monoclonal BOT-BNUB0936-500

Article Name	Thomsen-Friedenreich Antigen / CD176 (Pan Carcinoma Marker)(A78-G/A7), 0.2mg/mL, Clone: [A78-G/A7], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB0936-500
Supplier Catalog Number	BNUB0936-500
Alternative Catalog Number	BOT-BNUB0936-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human, Mouse, Rat
Immunogen	Neuraminidase-treated human red blood cells
Product Description	This antibody recognizes a disaccharide epitope, Gal 1-3GalNAc, of Thomsen-Friedenreich (TF) antigen. It is specific for both anomeric forms of the disaccharide (TF and TF , including related structures on the glycolipid) and shows no cross-reactivit...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[A78-G/A7]
Molecular Weight	Multiple

UniProt	Not Applicable
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Immunohistochemistry (formalin-fixed): 0.5-1.0 ug/mL for 30 minutes at RT Immunofluorescence: 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user