

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Mucin 3(M3.1), 0.2mg/mL, Clone: [M3.1], Mouse, Monoclonal BOT-BNUB0990-100**

Article Name	Mucin 3(M3.1), 0.2mg/mL, Clone: [M3.1], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB0990-100
Supplier Catalog Number	BNUB0990-100
Alternative Catalog Number	BOT-BNUB0990-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	A synthetic peptide of 35 amino acids, SIB35 (C-HSTPSFTSSITTTETTSHTSTPSFTSSITTTETTS), which contains two of the MUC3 tandem repeats, coupled to KLH.
Product Description	This antibody recognizes a protein of HMW, identified as mucin 3 glycoprotein (MUC3). Its epitope localizes between aa SITTTE. This MAb shows no cross-reaction with human milk fat globule membranes, MUC1, or MUC2. MUC3 is distributed in colon and rec...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[M3.1]
Molecular Weight	1100 kDa

UniProt	<a href="#">Q02505</a>
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user