

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Lewis B(2-25LE), 0.2mg/mL, Clone: [2-25LE], Mouse, Monoclonal BOT-BNUB1008-500

| | |
|----------------------------|---|
| Article Name | Lewis B(2-25LE), 0.2mg/mL, Clone: [2-25LE], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNUB1008-500 |
| Supplier Catalog Number | BNUB1008-500 |
| Alternative Catalog Number | BOT-BNUB1008-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Guinea pig, Human, Porcine |
| Immunogen | Mucin isolated from a human ovarian cyst fluid |
| Product Description | The Lewis histo-blood group system comprises a set of fucosylated glycosphingolipids that are synthesized by exocrine epithelial cells and circulate in body fluids. The glycosphingolipids function in embryogenesis, tissue differentiation, tumor metas... |
| Clonality | Monoclonal |
| Concentration | 0.2 mg/mL |
| Clone Designation | [2-25LE] |
| Molecular Weight | 45 kDa |
| UniProt | P21217 |

| | |
|-------------------|--|
| Buffer | PBS, 0.05% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user</p> |