

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

bcl-2(BCL2/782 + BCL2/796), 0.2mg/mL, Clone: [BCL2/782 BCL2/796], Mouse, Monoclonal BOT-BNUB1023-100

| | |
|----------------------------|---|
| Article Name | bcl-2(BCL2/782 + BCL2/796), 0.2mg/mL, Clone: [BCL2/782 BCL2/796], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNUB1023-100 |
| Supplier Catalog Number | BNUB1023-100 |
| Alternative Catalog Number | BOT-BNUB1023-100-100UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human bcl-2 protein |
| Product Description | This antibody recognizes a protein of 25-26 kDa, identified as Bcl-2. This antibody shows no cross-reaction with Bcl-x or Bax proteins. Bcl-2 suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural ... |
| Clonality | Monoclonal |
| Concentration | 0.2 mg/mL |
| Clone Designation | [BCL2/782 BCL2/796] |
| Molecular Weight | 25-26 kDa |

| | |
|-------------------|--|
| UniProt | P10415 |
| Buffer | PBS, 0.05% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |