

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Helicobacter pylori(B-53.4), 0.2mg/mL, Clone: [B-53.4], Mouse, Monoclonal BOT-BNUB1024-500

| | |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Article Name | Helicobacter pylori(B-53.4), 0.2mg/mL, Clone: [B-53.4], Mouse, Monoclonal |
| Biozol Catalog Number | BOT-BNUB1024-500 |
| Supplier Catalog Number | BNUB1024-500 |
| Alternative Catalog Number | BOT-BNUB1024-500-500UL |
| Manufacturer | Biotium |
| Host | Mouse |
| Category | Antikörper |
| Species Reactivity | H. pylori |
| Immunogen | Total sonicate of Helicobacter pylori |
| Product Description | The spiral shaped bacterium Helicobacter pylori is strongly associated with inflammation of the stomach and is also implicated in the development of gastric malignancy. H. pylori is known to cause peptic ulcers and chronic gastritis in human. It is assoc... |
| Clonality | Monoclonal |
| Concentration | 0.2 mg/mL |
| Clone Designation | [B-53.4] |
| Molecular Weight | Not Known |
| UniProt | Not Known |
| Buffer | PBS, 0.05% BSA, 0.05% azide |

| Source | Animal |
|-------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Application Notes | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |