

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **S100B(4C4.9 + S100B/1012), 0.2mg/mL, Clone: [4C4.9 S100B/1012], Mouse, Monoclonal BOT-BNUB1204-500**

|                            |   |
|----------------------------|---|
| Article Name               | S100B(4C4.9 + S100B/1012), 0.2mg/mL, Clone: [4C4.9 S100B/1012], Mouse, Monoclonal   |
| Biozol Catalog Number      | BOT-BNUB1204-500  |
| Supplier Catalog Number    | BNUB1204-500  |
| Alternative Catalog Number | BOT-BNUB1204-500-500UL  |
| Manufacturer               | Biotium   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | IF, IHC   |
| Species Reactivity         | Bovine, Human, Mouse, Rat   |
| Immunogen                  | Purified bovine brain S100 protein (4C4.9), Recombinant full-length human S100B protein (S100B/1012)  |
| Product Description        | S100 belongs to the family of calcium binding proteins. S100A and S100B proteins are two members of the S100 family. S100A is composed of an alpha and a beta chain whereas S100B is composed of two beta chains. This antibody is specific against an epi... |
| Clonality                  | Monoclonal  |
| Concentration              | 0.2 mg/mL   |
| Clone Designation          | [4C4.9 S100B/1012]  |
| Molecular Weight           | 10-12 kDa   |

|                   |   |
|-------------------|---|
| UniProt           | <a href="#">P04271</a>  |
| Buffer            | PBS, 0.05% BSA, 0.05% azide   |
| Source            | Animal  |
| Application Notes | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p> |