

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **TDP2 / EAPII (TDP2/1258), 0.2mg/mL, Clone: [TDP2/1258], Mouse, Monoclonal BOT-BNUB1258-500**

Article Name	TDP2 / EAPII (TDP2/1258), 0.2mg/mL, Clone: [TDP2/1258], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB1258-500
Supplier Catalog Number	BNUB1258-500
Alternative Catalog Number	BOT-BNUB1258-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant human full-length TDP2 protein
Product Description	This MAb recognizes a protein of 41 kDa, which is identified as TDP2. It is a member of a superfamily of divalent cation-dependent phosphodiesterases. The encoded protein associates with CD40, tumor necrosis factor (TNF) receptor-75 and TNF receptor ...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[TDP2/1258]
Molecular Weight	~41 kDa

UniProt	<a href="#">O95551</a>
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user