

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

p57Kip2 (Mitotic Inhibitor/Suppressor Protein)(57P06), 0.2mg/mL, Clone: [57P06], Mouse, Monoclonal BOT-BNUB1264-500

Article Name	p57Kip2 (Mitotic Inhibitor/Suppressor Protein)(57P06), 0.2mg/mL, Clone: [57P06], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB1264-500
Supplier Catalog Number	BNUB1264-500
Alternative Catalog Number	BOT-BNUB1264-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human, Mouse
Immunogen	Recombinant full-length human p57Kip2 protein
Product Description	Recognizes a protein of 57 kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as a...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[57P06]
Molecular Weight	57 kDa

UniProt	P49918
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 0.25-0.5 ug/mL for 30 minutes at RT Flow cytometry: 0.5-1 ug/million cells Immunofluorescence: 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user