

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Villin (GI-Mucosal & Urogenital Brush Border Marker) (VIL1/1314), 0.2mg/mL, Clone: [VIL1/1314], Mouse, Monoclonal BOT-BNUB1314-500

Article Name	Villin (GI-Mucosal & Urogenital Brush Border Marker) (VIL1/1314), 0.2mg/mL, Clone: [VIL1/1314], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB1314-500
Supplier Catalog Number	BNUB1314-500
Alternative Catalog Number	BOT-BNUB1314-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC, WB
Species Reactivity	Human
Immunogen	Recombinant human Villin fragment of 133 amino acid residues (aa179-311) (exact sequence is proprietary)
Product Description	This antibody recognizes a protein of 95 kDa, which is identified as villin. It is a major constituent in the microvilli, which compose the brush border of epithelial cells forming absorptive surfaces of the intestinal and renal proximal tubular epit...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[VIL1/1314]
Molecular Weight	93 kDa

UniProt	P09327
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Flow cytometry: 0.5-1 ug/million cells in 0.1mL Immunohistology (formalin): 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Western blotting 1-2 ug/mL Optimal dilution for a specific application should be determined by user