

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **CD31 / PECAM-1 (Endothelial Cell Marker) (C31/1395R), 0.2mg/mL, Clone: [C31/1395R], Rabbit, Monoclonal BOT-BNUB1395-500**

Article Name	CD31 / PECAM-1 (Endothelial Cell Marker) (C31/1395R), 0.2mg/mL, Clone: [C31/1395R], Rabbit, Monoclonal
Biozol Catalog Number	BOT-BNUB1395-500
Supplier Catalog Number	BNUB1395-500
Alternative Catalog Number	BOT-BNUB1395-500-500UL
Manufacturer	Biotium
Host	Rabbit
Category	Antikörper
Application	FC, IHC, WB
Species Reactivity	Human
Immunogen	Human recombinant full-length CD31 protein
Product Description	CD31 (PECAM-1) is a transmembrane glycoprotein member of the immunoglobulin supergene family of adhesion molecules. CD31 is expressed by stem cells of the hematopoietic system and is primarily used to identify and concentrate these cells for experime...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[C31/1395R]
Molecular Weight	~100 kDa (endothelium) ~130 kDa (platelets)

UniProt	<a href="#">P16284</a>
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user