

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

MART-1 / Melan-A / MLANA (Melanoma Marker) (MLANA/1409R), 0.2mg/mL, Clone: [MLANA/1409R], Rabbit, Monoclonal BOT-BNUB1409-500

| | |
|----------------------------|---|
| Article Name | MART-1 / Melan-A / MLANA (Melanoma Marker) (MLANA/1409R), 0.2mg/mL, Clone: [MLANA/1409R], Rabbit, Monoclonal |
| Biozol Catalog Number | BOT-BNUB1409-500 |
| Supplier Catalog Number | BNUB1409-500 |
| Alternative Catalog Number | BOT-BNUB1409-500-500UL |
| Manufacturer | Biotium |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human, Mouse, Rat |
| Immunogen | Recombinant human full-length MLANA protein |
| Product Description | This antibody recognizes a protein doublet of 20-22 kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Sev... |
| Clonality | Monoclonal |
| Concentration | 0.2 mg/mL |
| Clone Designation | [MLANA/1409R] |
| Molecular Weight | 20-22 kDa (doublet) |

| | |
|-------------------|--|
| UniProt | Q16655 |
| Buffer | PBS, 0.05% BSA, 0.05% azide |
| Source | Animal |
| Application Notes | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |