

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Lewis Y (Tumor Marker) (A70-A/A9), 0.2mg/mL, Clone: [A70-A/A9], Mouse, Monoclonal BOT-BNUB1412-100

Article Name	Lewis Y (Tumor Marker) (A70-A/A9), 0.2mg/mL, Clone: [A70-A/A9], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB1412-100
Supplier Catalog Number	BNUB1412-100
Alternative Catalog Number	BOT-BNUB1412-100-100UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Live Ls174T cells (human colon carcinoma cell line)
Product Description	This antibody recognizes a carbohydrate epitope present on tumor-associated Lewis Y antigen (Fucal α 1-2Gal β 1-4/3GlcNAc β). Lewis Y is expressed in large bowel tumors and colorectal carcinomas. It may be useful in the classification of human ren...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[A70-A/A9]
Molecular Weight	Multiple

UniProt	Not Applicable
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user