

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD3e (T-Cell Marker) (PC3/188A), 0.2mg/mL, IgG1, Clone: [PC3/188A], Mouse, Monoclonal BOT-BNUB1479-500

Article Name	CD3e (T-Cell Marker) (PC3/188A), 0.2mg/mL, IgG1, Clone: [PC3/188A], Mouse, Monoclonal
Biozol Catalog Number	BOT-BNUB1479-500
Supplier Catalog Number	BNUB1479-500
Alternative Catalog Number	BOT-BNUB1479-500-500UL
Manufacturer	Biotium
Host	Mouse
Category	Antikörper
Application	IHC, WB
Species Reactivity	Human
Immunogen	Synthetic peptide corresponding to amino acids 156-168 (ERPPPVPNPDYEP) of the cytoplasmic domain of human CD3e
Product Description	The immunogen for this Mab is derived from the cytoplasmic domain of CD3 epsilon chain. CD3 consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28 kDa. The CD3 complex is closely ...
Clonality	Monoclonal
Concentration	0.2 mg/mL
Clone Designation	[PC3/188A]
Molecular Weight	20 kDa

Isotype	IgG1
UniProt	P07766
Buffer	PBS, 0.05% BSA, 0.05% azide
Source	Animal
Application Notes	Not recommended for flow cytometry Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Western blotting 1-2 ug/mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user